arXiv:math-ph/0608015v1 4 Aug 2006

STURM-LIOUVILLE PROBLEM IN QUANTUM CALCULUS

AHMED FITOUHI, AKRAM NEMRI, AND MENIAR HADDAD

Abstract. This paper aims to study the q-analogue of the Sturm Liouville problem and to give an asymptotic behaviour at infinity for its solution φ. Additionally, we establish an asymptotic expansion of the q-Bessel function j_α for $\alpha > -\frac{1}{2}$. We are not in situation to claim that our results are new but they have the advantage to show that the method used by Agranovich and Marchenko remain true.

1. Introduction:

In the classical spectral analysis (see [2] , [14],...), we denote by L a linear differential operator of the second-order given by on $[0,\infty[$ of the form

$$Lu = \frac{d^2u}{dx^2} - p(x)u \tag{1.1}$$

where $p(x)$ is a real function, continuous which is integrable on $[0,\infty[$.

We note by $\varphi(x,\lambda^2)$ and $\theta(x,\lambda^2)$ the solutions of

$$Lu = \lambda^2 u \tag{1.2}$$

with initial conditions

$$\begin{array}{ll} \varphi(0,\lambda^2) = \sin\alpha, & \varphi'(0,\lambda^2) = -\cos\alpha; \\ \theta(0,\lambda^2) = \cos\alpha, & \theta'(0,\lambda^2) = \sin\alpha, \end{array}$$

where λ is an arbitrary positif real number and α is an arbitrary real number.

It is known that solving (1.2) is equivalent to solving the following volterra integral equation

$$u(x,\lambda^2) = \sin\alpha\cos(\lambda x) - \frac{\sin(\lambda x)}{\lambda}\cos\alpha + \int_0^x \frac{\sin(\lambda(x-y))}{\lambda} p(y)u(y,\lambda^2)dy \tag{1.3}$$

where $x \in [0,+\infty[$, $\lambda \in \mathbb{R}_+^*$ and $p(x)$ is an continuous integrable function on $[0,+\infty[$. Hence, for all $\lambda \geq \rho > 0$, $\varphi(x,\lambda^2)$ is a bounded function and have the asymptotic formulas

$$\varphi(x,\lambda^2) = \mu(\lambda^2)\cos(\lambda x) + \nu(\lambda^2)\sin(\lambda x) + O(1) \tag{1.4}$$

where

$$\mu(\lambda^2) = \sin\alpha - \int_0^\infty \frac{\sin(\lambda y)}{\lambda} p(y)\varphi(y,\lambda^2)dy, \tag{1.5}$$

$$\nu(\lambda^2) = -\frac{\cos\alpha}{\lambda} + \int_0^\infty \frac{\cos(\lambda y)}{\lambda} p(y)\varphi(y,\lambda^2)dy. \tag{1.6}$$

2000 *Mathematics Subject Classification.* 33D60, 26D15, 33D05, 33D15, 33D90.

Key words and phrases. Quantum calculus, q-analysis, q-Integral Transform.

Similarly, we have

$$\theta(x,\lambda^2)=\mu_1(\lambda^2)\cos(\lambda x)+\nu_1(\lambda^2)\sin(\lambda x)+O(1) \tag{1.7}$$

where

$$\mu_1(\lambda^2) = \cos\alpha-\int_0^\infty\frac{\sin(\lambda y)}{\lambda}p(y)\theta(y,\lambda^2)dy, \tag{1.8}$$

$$\nu_1(\lambda^2) = \frac{\sin\alpha}{\lambda}+\int_0^\infty\frac{\cos(\lambda y)}{\lambda}p(y)\theta(y,\lambda^2)dy. \tag{1.9}$$

In the present paper we are concerned to give its q-analogue and study its asymptotic behaviour at infinity.
This paper is organized as follows: in section 2, we present some preliminaries results and notations that will be useful in the sequel. Further it is natural to consider in section 3, the asymptotic behaviour of $\varphi(x,\lambda^2;q^2)$ and $\theta(x,\lambda^2)$ for $\lambda\longrightarrow\infty$. the fundamental result is given in the following theorem

Theorem 1.1. *For λ in $\mathbb{R}_{q,+}$, we have:*

$$\mu(\lambda^2;q^2)\nu_1(\lambda^2;q^2)-\nu(\lambda^2;q^2)\mu_1(\lambda^2;q^2)=\frac{1}{q^{\frac{1}{2}}\lambda}. \tag{1.10}$$

Section 4, is devoted to finding precise asymptotic formulas of j_α: the q-Bessel function for large λ.

2. Notations and preliminaries

We recall some usual notions and notations used in the q-theory. Let a and q be real numbers such that $0<q<1$. In all the sequel we suppose that and $\dfrac{\log(1-q)}{\log q}\in\mathbb{Z}$.
The q-shifted factorials are defined by

$$(a;q)_n = \prod_{k=0}^{n-1}(1-aq^k) \quad ;n\in\mathbb{N}\backslash\{0\}, \tag{2.1}$$

$$(a;q)_0 = 1, \tag{2.2}$$

$$(a;q)_\infty = \prod_{k=0}^{\infty}(1-aq^k) \tag{2.3}$$

and more generally:

$$(a_1,\cdots,a_r;q)_n = \prod_{k=1}^{r}(a_k;q)_n. \tag{2.4}$$

The basic hypergeometric series or q-hypergeometric series is given for r , s integers by

$$_r\phi_s(a_1,\cdots,a_r;b_1,\cdots,b_s;q,x)=\sum_{n=0}^{\infty}\frac{(a_1,\cdots,a_r;q)_n}{(b_1,\cdots,b_s;q)_n(q,q)_n}[(-1)^nq^{\frac{n(n-1)}{2}}]^{1+s-r}x^n$$

here

$$r,s\in\mathbb{N};a_1,\cdots,a_r\in\mathbb{C};b_1,\cdots,b_s\in\mathbb{C}\backslash\{q^{-k},k\in\mathbb{N}\}$$

The q-derivative $D_{q,x}f$ of a function f on an open interval is given by:

$$D_{q,x}f(x) = \frac{f(x) - f(qx)}{(1-q)x}, \quad x \neq 0 \tag{2.5}$$

and $(D_{q,x}f)(0) = f^{'}(0)$ provided $f^{'}(0)$ exist. The q-shift operators are

$$(\Lambda_{q,x}f)(x) = f(qx) \tag{2.6}$$

$$(\Lambda_{q,x}^{-1}f)(x) = f(q^{-1}x). \tag{2.7}$$

We consider the q-operator

$$\Delta_{q,x} = \Lambda_{q,x}^{-1} D_{q,x}^2. \tag{2.8}$$

The q-Jackson integral from 0 to a and from 0 to ∞ are respectively defined by

$$\int_0^a f(x)d_qx = (1-q)a\sum_{n=0}^{\infty} f(aq^n)q^n \tag{2.9}$$

$$\int_0^{\infty} f(x)d_qx = (1-q)\sum_{-\infty}^{+\infty} f(q^n)q^n \tag{2.10}$$

and from a to ∞,

$$\int_a^{\infty} f(x)d_qx = (1-q)a\sum_{n=1}^{+\infty} f(aq^{-n})q^{-n}. \tag{2.11}$$

Some q-functional spaces will be used to establish our result. We begin by putting

$$\mathbb{R}_q = \{\pm q^k, k \in \mathbb{Z}\} \cup \{0\}, \tag{2.12}$$

$$\mathbb{R}_{q,+} = \{+q^k, k \in \mathbb{Z}\}. \tag{2.13}$$

Let $L_q^p(\mathbb{R}_{q,+})$, $p \in [1, +\infty[$ be the space of functions f such that

$$\parallel f \parallel_{q,p} = (\int_0^{\infty} \mid f(x) \mid^p d_qx)^{\frac{1}{p}} < +\infty, \tag{2.14}$$

and for $p = \infty$

$$\parallel f \parallel_{q,\infty} = ess \sup_{x \in \mathbb{R}_{q,+}} \mid f(x) \mid \tag{2.15}$$

Note that for $n \in \mathbb{Z}$ and $a \in \mathbb{R}_q$, we have

$$\int_0^{\infty} f(q^nx)d_qx = \frac{1}{q^n}\int_0^{\infty} f(x)d_qx. \tag{2.16}$$

$$\int_0^{a} f(q^nx)d_qx = \frac{1}{q^n}\int_0^{aq^n} f(x)d_qx. \tag{2.17}$$

The q-integration by parts is given for suitable function f and g by:

$$\int_0^{\infty} f(x)D_{q,x}g(x)d_qx = \Big[f(x)g(x)\Big]_0^{\infty} - \int_0^{\infty} D_{q,x}(f(q^{-1}x))g(x)d_qx. \tag{2.18}$$

Jackson in [8] defined the q-analogue of the Gamma function as

$$\Gamma_q(x) = \frac{(q;q)_{\infty}}{(q^x;q)_{\infty}}(1-q)^{1-x} \tag{2.19}$$

We take the definition of q-trigonometric given by T.H.Koornwinder and R.F.Swarttouw (see [11]) with simple changes and we write q-cosine and q-sinus as a series of functions

$$\cos(x;q^2) = {}_1\varphi_1(0,q,q^2;(1-q)^2x^2) = \sum_{n=0}^{\infty}(-1)^n b_n(x;q^2) \tag{2.20}$$

$$\sin(x;q^2) = (1-q)x{}_1\varphi_1(0,q^3,q^2;(1-q)^2x^2) = \sum_{n=0}^{\infty}(-1)^n c_n(x;q^2) \tag{2.21}$$

where we have put

$$b_n(x;q^2) = b_n(1;q^2)x^{2n} = q^{n(n-1)}\frac{(1-q)^{2n}}{(q;q)_{2n}}x^{2n} \tag{2.22}$$

$$c_n(x;q^2) = c_n(1;q^2)x^{2n+1} = q^{n(n-1)}\frac{(1-q)^{2n+1}}{(q;q)_{2n+1}}x^{2n+1}. \tag{2.23}$$

The reader will notice that the previous definition (2.20) derived from those given in [11] with minor change. These functions satisfy

$$D_{q,x}\cos(x;q^2) = -q^{-1}\sin(qx;q^2), \tag{2.24}$$

$$D_{q,x}\sin(x;q^2) = \cos(x;q^2). \tag{2.25}$$

and we have the following estimations:

$$|\cos(x;q^2)| \leq \frac{1}{(q;q^2)^2_\infty}, \tag{2.26}$$

$$|\sin(x;q^2)| \leq \frac{1}{(q;q^2)^2_\infty}. \tag{2.27}$$

We recall the tow q-analogue of the exponential functions [10], defined by:

$$E(x;q) = (-(1-q)x;q)_\infty = \sum_{n=0}^{\infty}\frac{(1-q)^n}{(q;q)_n}q^{n(n-1)/2}x^n, \qquad x \in \mathbb{R} \tag{2.28}$$

$$e(x;q) = \frac{1}{((1-q)x;q)_\infty} = \sum_{n=0}^{\infty}\frac{(1-q)^n}{(q;q)_n}x^n, \qquad |x| < \frac{1}{1-q}. \tag{2.29}$$

The function $E(x;q)$ is analytic and $e(z;q)$ is a meromorphic function on $\mathbb{C}$ having simple poles at $z = \frac{q^{-m}}{1-q}$, $m \in \mathbb{N}$. They satisfy

$$e(x;q)E(-x;q) = 1. \tag{2.30}$$

Proposition 2.1. *1- If F is any q-derivative of the function f, namely $D_{q,x}F(x) = f(x)$, continuous at $x = 0$, then*

$$\int_0^x f(t)d_qt = (F(x) - F(0)). \tag{2.31}$$

2- For any function f we have

$$D_{q,x}\left[\int_0^x f(t)d_qt\right] = f(x) \tag{2.32}$$

3- If G is any q-derivative of the function g, integrable over (x,∞); $x \geq 0$ then:

$$\int_x^{+\infty} g(t)d_qt = -\lim_{b\longrightarrow+\infty}(G(x)-G(b)) \quad = \quad G(\infty)-G(x) \tag{2.33}$$

4- For any function f integrable over $(x,+\infty)$, we have

$$D_{q,x}\left[\int_x^{+\infty} g(t)d_qt\right] = -g(x) \tag{2.34}$$

2.1. **The q-Wronskian:** Let the following q-difference equation:

$$D^2_{q,x}u(x)+a(x)D_{q,x}u(x)+b(x)u(qx)=0 \tag{2.35}$$

Proposition 2.2. *We define the q-Wronskian $W(x;q)$ For the two solutions u_1 and u_2 of the q-difference equation (2.35) by:*

$$W(x;q)(x) \quad = \quad u_1(qx)D_{q,x}u_2(x)-u_2(qx)D_{q,x}u_1(x) \tag{2.36}$$

$$= \quad \frac{u_1(qx)u_2(x)-u_1(x)u_2(qx)}{(1-q)x}. \tag{2.37}$$

It satisfies the following q-difference equation:

$$D_{q,x}W(x;q)+a(x)W(x;q)=0. \tag{2.38}$$

Furthermore:

$$W(x;q)=\frac{W(0;q)}{\prod_{k=0}^{\infty}[1+(1-q)q^kxa(q^kx)]} \tag{2.39}$$

Proof. It easy to see that, if u_1 and u_2 are tow solutions of (2.35), we have the relation (2.38).
Using the q-derivative definition (2.5), the equation (2.38) can be rewritten as:

$$W(x;q)-W(qx;q)=-(1-q)xa(x)W(x;q) \tag{2.40}$$

then

$$W(qx,q)=\frac{1}{1+(1-q)qxa(qx)}W(q^2x;q). \tag{2.41}$$

So, by induction we have for $n\in\mathbb{N}$:

$$W(q^nx;q)=\frac{1}{1+(1-q)q^nxa(q^nx)}W(q^{n+1}x;q). \tag{2.42}$$

We deduce that

$$W(x;q)=\frac{W(0;q)}{\prod_{k=0}^{\infty}[1+(1-q)q^kxa(q^kx)]}. \tag{2.43}$$

$\square$

Proposition 2.3. *The solution of the q-difference equation*

$$(E) = \begin{cases} D_{q,x}^2 u(x) + u(qx) = 0 \\ u(0;q^2) = a \\ D_q u(0;q^2) = b \end{cases} \tag{2.44}$$

is given by:

$$u(x;q^2) = a\cos(x;q^2) + bq^{-\frac{1}{2}} sin(q^{\frac{1}{2}}x;q^2) \tag{2.45}$$

Proof. Let $u(x) = \sum_{n\geq 0} a_n x^n$. Then

$$D_{q,x}^2 u(x) = \sum_{n\geq 0} a_{n+2} \frac{1-q^{n+2}}{1-q}\frac{1-q^{n+1}}{1-q} x^n. \tag{2.46}$$

If we replace in (2.44) we have the following recurrence relation:

$$\frac{1-q^{n+2}}{1-q}\frac{1-q^{n+1}}{1-q} a_{n+2} = q^n a_n. \tag{2.47}$$

then

If $n = 2p$; $p \in \mathbb{N}$, we have

$$q^{2p} a_{2p} = \frac{1-q^{2p+2}}{1-q}\frac{1-q^{2p+1}}{1-q} a_{2p+2} \tag{2.48}$$

so, by induction on p, we obtain:

$$a_{2p} = (-1)^p \frac{(1-q)^{2p}}{(q;q)_{2p}} q^{p(p-1)} a_0. \tag{2.49}$$

similarly

if $n = 2p+1$, we obtain

$$a_{2p+1} = (-1)^p \frac{(1-q)^{2p+1}}{(q;q)_{2p+1}} q^{p(p-1)} a_1. \tag{2.50}$$

Using the definition (2.20), the solution of (2.44) is given. □

Corollary 2.4. *For $x \in \mathbb{R}_q$ we have*

$$\cos(qx;q^2)\cos(q^{\frac{1}{2}}x;q^2) + q^{-\frac{3}{2}} sin(q^{\frac{3}{2}}x;q^2) sin(qx;q^2) = 1 \tag{2.51}$$

which tends to the classical trigonometric relation

$$cos^2(x) + sin^2(x) = 1 \tag{2.52}$$

when $q \to 1^-$

Proof. Using the relation (2.38) and the condition $a(x) = 0$, we have

$$D_{q,x} W(x;q) = 0. \tag{2.53}$$

So,

$$W(x;q) = \lim_{n\to+\infty} W(q^n x;q) = W(0;q).$$

$$\begin{aligned} W(0;q) &= \lim_{x\to 0} W(x;q) = \lim_{x\to 0}\Big[\cos(qx;q^2)\cos(q^{\frac{1}{2}}x;q^2) + q^{-\frac{3}{2}}\sin(q^{\frac{3}{2}}x;q^2)\sin(qx;q^2)\Big] \\ &= \lim_{x\to 0}\frac{1}{(1-q)x}\Big[q^{-\frac{1}{2}}\cos(x;q^2)\sin(q^{\frac{3}{2}}x;q^2) - q^{-\frac{1}{2}}\cos(qx;q^2)\sin(q^{\frac{1}{2}}x;q^2)\Big] \\ &= 1 \end{aligned}$$

The result follows. □

3. Asymptotic expansion of solutions at infinity

In this section, we try to study for $\lambda \longrightarrow \infty$ the asymptotic expansion of solution $u(x,\lambda^2;q^2)$ of L_q the q-difference operator defined by:

$$L_q u(x) = D^2_{q,x}u(x) - p(x)u(x); \qquad p(x)\in L^{\infty}(\mathbb{R}_{q,+})\cap L^1(\mathbb{R}_{q,+}). \tag{3.1}$$

In the next we try to resolve the following q-difference problem

$$L_q u(x) = -\lambda^2 u(qx), \qquad x\in\mathbb{R}_{q,+}, \lambda\in\mathbb{R}_{q,+}. \tag{3.2}$$

Proposition 3.1. (The q-Gronwall lemma:)
Let f and g be two positive functions, continuous at 0 and q-integrable over all finite interval of $[0,+\infty[$.
We suppose that

$$f(x) \le C_q + \int_0^x f(t)g((t)d_qt \tag{3.3}$$

where $C_q\in\mathbb{R}_{q,+}$.
Then

$$f(x) \le \frac{C_q}{\displaystyle\prod_{k=0}^{\infty}[1-(1-q)q^kxg(q^kx)]} \tag{3.4}$$

Proof. Let the following q-Jackson integral

$$y(x) = \int_0^x f(t)g(t)d_qt,$$

we have,

$$D_{q,x}y(x) = f(x)g(x) \le [C_q + y(x)]g(x)$$

then,

$$C_q + y(x) \le \frac{1}{1-(1-q)xg(x)}(C_q + y(qx)$$

and by induction on n, we deduce that

$$C_q + y(q^nx) \le \frac{1}{1-(1-q)q^nxg(q^nx)}(C_q + y(q^{n+1}x))$$

then

$$C_q + y(x) \le \frac{1}{\displaystyle\prod_{k=0}^{\infty}[1-(1-q)q^kxg(q^kx)]}(C_q + y(0)).$$

The fact that $y(0)=0$ leads to the result. □

Corollary 3.2. *Let f be a positive function, continuous at 0 and q-integrable over all finite interval of* $[0,+\infty[$. *We suppose that there exist two constants C_q and M_q in $\mathbb{R}_{q,+}$, such that*

$$f(x) \leq C_q + M_q \int_0^x f(t) d_q t.$$

Then we have,

$$f(x) \leq C_q e(M_q(1-q)x; q^2) \tag{3.5}$$

where $e(x;q^2)$ is given by (2.29).

Definition 3.3. Let $U(x;q^2)$ and $V(x;q^2)$ be twice q-differentiable functions. We define $[U,V]_q$ by:

$$[U,V]_q = U(qx;q^2) D_{q,x} V(x;q^2) - V(qx;q^2) D_{q,x} U(x;q^2) \tag{3.6}$$

Proposition 3.4. *(q-Green formula)*
For $U(x;q^2)$ and $V(x;q^2)$ twice q-differentiable functions, we have

$$D_{q,x} [U,V]_q = V(qx;q^2) L_q U(x;q^2) - U(qx;q^2) L_q V(x;q^2) \tag{3.7}$$

3.1. **q-Asymptotic behaviour of $\varphi(x,\lambda^2;q^2)$ when $\lambda \longrightarrow \infty$.** For $\lambda \in \mathbb{R}_{q,+}$ let $\varphi(x,\lambda^2;q^2)$ the solution of the following q-difference problem (E_1)

$$(E_1) = \begin{cases} L_q U(x,\lambda^2;q^2) = -\lambda^2 U(qx,\lambda^2;q^2), \\ U(0,\lambda^2;q^2) = q^{-1} \sin(q\alpha;q^2), \\ D_q U(0,\lambda^2;q^2) = \cos(q\alpha;q^2) \quad , \alpha \in \mathbb{R}. \end{cases} \tag{3.8}$$

where L_q is given by (3.2).

Theorem 3.5. *Let $p(x)$ in $L_q^\infty(\mathbb{R}_{q,+})$, then the solution $\varphi(x,\lambda^2;q^2)$ of (E_1) verifies the following q-integral equation:*

$$\begin{aligned} \varphi(x,\lambda^2;q^2) &= q^{-1} \sin(q\alpha;q^2) \cos(\lambda x;q^2) + q^{-\frac{1}{2}} \frac{\cos(q\alpha;q^2)}{\lambda} \sin(q^{\frac{1}{2}}\lambda x;q^2) \\ &+ \frac{1}{\lambda} \int_0^x G(x,y,\lambda^2;q^2) p(y) \varphi(y,\lambda^2;q^2) d_q y, \end{aligned} \tag{3.9}$$

where $G(x,y,\lambda^2;q^2)$ is the Green kernel defined by

$$G(x,y,\lambda^2;q^2) = \cos(q\lambda y;q^2) sin(q^{\frac{1}{2}}\lambda x;q^2) - sin(q^{\frac{3}{2}}\lambda y;q^2) \cos(\lambda x;q^2) \tag{3.10}$$

Proof. We begin by resolving the following q-homogenous equation $(E_{1,h})$

$$(E_{1,h}): \qquad D^2_{q,x} U(x,\lambda^2;q^2) + \lambda^2 U(qx,\lambda^2;q^2) = 0.$$

For this way, if we use the same steps, given in proposition 2.3, we obtain

$$\varphi_{1,h}(x,\lambda^2;q^2) = a\cos(\lambda x;q^2) + b\frac{q^{-\frac{1}{2}}}{\lambda} \sin(q^{\frac{1}{2}}\lambda x;q^2) \quad ; a,b \in \mathbb{R}.$$

Now we are able to give a particular solution $\varphi_p(x,\lambda^2;q^2)$ of (E_1) . For deeps, we use the q-Method of variation of constant. Hence we write $\varphi_p(x,\lambda^2;q^2)$ in the following form

$$\varphi_p(x,\lambda^2;q^2) = a(x,\lambda^2;q^2)\cos(\lambda x;q^2) + b(x,\lambda^2;q^2)\frac{q^{-\frac{1}{2}}}{\lambda} \sin(q^{\frac{1}{2}}\lambda x;q^2)$$

Therefore, if we replace $D_{q,x}\varphi(x,\lambda^2;q^2)$ and $D^2_{q,x}\varphi(x,\lambda^2;q^2)$ in (E_1), it can be rewritten in the form

$$D_{q,x}\left[I_1\right] + I_2 = p(x)\varphi(x,\lambda^2;q^2), \tag{3.11}$$

where

$$I_1 = D_{q,x}a(x,\lambda^2;q^2)\cos(q\lambda x;q^2) + D_{q,x}b(x,\lambda^2;q^2)\sin(q^{\frac{3}{2}}\lambda x;q^2) \tag{3.12}$$

and

$$I_2 = D_{q,x}a(x,\lambda^2;q^2)D_{q,x}[\cos(\lambda x;q^2)] + D_{q,x}b(x,\lambda^2;q^2)D_{q,x}[\sin(q^{\frac{1}{2}}\lambda x;q^2)]. \tag{3.13}$$

On the other hand, if we use (2.24) and (2.25), we obtain the following system

$$\left\{\begin{array}{lcl} D_{q,x}a(x,\lambda^2;q^2)\cos(q\lambda x;q^2) + D_{q,x}b(x,\lambda^2;q^2)sin(q^{\frac{3}{2}}\lambda x;q^2) & = & 0 \\ D_{q,x}a(x,\lambda^2;q^2)D_{q,x}[\cos(\lambda x;q^2)] + D_{q,x}b(x,\lambda^2;q^2)D_{q,x}[\sin(q^{\frac{1}{2}}\lambda x;q^2)] & = & p(x)\varphi(x,\lambda^2;q^2) \end{array}\right. \tag{3.14}$$

Hence ,

$$\begin{aligned} D_{q,x}a(x,\lambda^2;q^2) &= \frac{1}{W(x,\lambda^2;q^2)}\begin{vmatrix} 0 & sin(q^{\frac{3}{2}}\lambda x;q^2) \\ p(x)\varphi(x,\lambda^2;q^2) & q^{\frac{1}{2}}\lambda\cos(q^{\frac{1}{2}}\lambda x;q^2) \end{vmatrix} \\ &= -\frac{p(x)\varphi(x,\lambda^2;q^2)\sin(q^{\frac{3}{2}}\lambda x;q^2)}{W(x,\lambda^2;q^2)}, \end{aligned}$$

where $W(x,\lambda^2;q^2)$ is given by (2.36). So by proposition 2.1 and proposition 2.3 we obtain respectively that

$$W(x,\lambda^2;q^2) = \lambda \tag{3.15}$$

and

$$a(x,\lambda^2;q^2) = -\frac{1}{\lambda}\int_0^x p(y)\varphi(y,\lambda^2;q^2)\sin(q^{\frac{3}{2}}\lambda y;q^2)d_qy. \tag{3.16}$$

In a similar way, we can show that

$$b(x,\lambda^2;q^2) = \frac{1}{\lambda}\int_0^x p(y)\varphi(y,\lambda^2;q^2)\cos(q\lambda y;q^2)d_qy \tag{3.17}$$

then the particular solution $\varphi_p(x,\lambda^2;q^2)$ of (E_1) is given by

$$\varphi_p(x,\lambda^2;q^2) = \frac{1}{\lambda}\int_0^x p(y)\varphi(y,\lambda^2;q^2)G(x,y,\lambda^2;q^2)d_qy. \tag{3.18}$$

where

$$G(x,y,\lambda^2;q^2) = cos(q\lambda y;q^2)sin(q^{\frac{1}{2}}\lambda x;q^2) - sin(q^{\frac{3}{2}}\lambda y;q^2)\cos(q\lambda x;q^2) \tag{3.19}$$

furthermore, by the fact that

$$\varphi(0,\lambda^2) = a = q^{-1}sin(q\alpha;q^2)$$

and

$$D_q\varphi(0;q^2) = b = cos(q\alpha;q^2)$$

we can deduce the result. □

Proposition 3.6. *Let* $p(x)$ *a boundary function on* $\mathbb{R}_{q,+}$. *Then* $\varphi(x,\lambda^2;q^2)$ *verifies:*

(1) *For* $\lambda \in \mathbb{R}_{q,+}$,

$$\varphi(x,\lambda^2;q^2) = \mathcal{O}\left(e(C_\lambda(1-q)x;q^2)\right) \tag{3.20}$$

where

$$C_\lambda = \frac{2 \parallel p \parallel_{q,\infty}}{\mid \lambda \mid (q;q^2)^2_\infty} \tag{3.21}$$

and $e(x;q^2)$ *is given by (2.29).*

(2) *Additionally, if* $\lambda \longrightarrow \infty$, *we have*

$$\varphi(x,\lambda^2;q^2) = \mathcal{O}(1;q^2) \tag{3.22}$$

Proof. To prove the first result, it suffices to use theorem 3.5, (2.26) and (2.27). Therefore,

$$\begin{aligned}
\mid \varphi(x,\lambda^2;q^2) \mid \quad &\leq \quad \frac{\mid a \mid}{(q;q^2)^2_\infty} + \frac{\mid b \mid}{q \mid \lambda \mid (q;q^2)^2_\infty} + \frac{2}{\mid \lambda \mid (q;q^2)^2_\infty} \int_0^x \mid \varphi(y,\lambda^2;q^2) \mid\mid p(y) \mid d_q y, \\
&\leq \quad \frac{\mid a \mid}{(q;q^2)^2_\infty} + \frac{\mid b \mid}{q|\lambda|(q;q^2)^2_\infty} + \frac{2}{\mid \lambda \mid (q;q^2)^2_\infty} \parallel p \parallel_{q,\infty} \int_0^x \mid \varphi(y,\lambda^2;q^2) \mid d_q y, \\
\mid \varphi(x,\lambda^2;q^2) \mid \quad &\leq \quad A_\lambda + C_\lambda \int_0^x \mid \varphi(y,\lambda^2;q^2) \mid d_q y,
\end{aligned}$$

where

$$A_\lambda = \frac{\mid a \mid}{(q;q^2)^2_\infty} + \frac{\mid b \mid}{q \mid \lambda \mid (q;q^2)^2_\infty}$$

and

$$C_\lambda = \frac{\parallel p \parallel_{q,\infty}}{\mid \lambda \mid (q;q^2)^2_\infty}$$

By proposition 3.2, we obtain that

$$\mid \varphi(x,\lambda^2;q^2) \mid \leq A_\lambda e(C_\lambda(1-q)x;q^2). \tag{3.23}$$

and the result follows immediately. □

Theorem 3.7. *For* $p(x)$ *in* $L^\infty_q(\mathbb{R}_{q,+}) \cap L^1_q(\mathbb{R}_{q,+})$, *we have*

(1) *For* $\lambda \geq \xi > 0$, $\varphi(x,\lambda^2;q^2)$ *is an uniformly bounded function.*

(2) *For a large* λ , *we have the following estimation*

$$\varphi(x,\lambda^2;q^2) = \mu(\lambda^2;q^2)\cos(\lambda x;q^2) + \nu(\lambda^2;q^2)q^{-\frac{1}{2}}\sin(q^{\frac{1}{2}}\lambda x;q^2) + \mathcal{O}(1;q^2) \tag{3.24}$$

where

$$\mu(\lambda^2;q^2) = q^{-1}\sin(q\alpha;q^2) - \frac{1}{\lambda}\int_0^\infty sin(q^{\frac{3}{2}}\lambda y;q^2)p(y)\varphi(y,\lambda^2;q^2)d_q y \tag{3.25}$$

and

$$\nu(\lambda^2;q^2) = \frac{\cos(q\alpha;q^2)}{\lambda} + \frac{q^{\frac{1}{2}}}{\lambda}\int_0^\infty \cos(q\lambda y;q^2)p(y)\varphi(y,\lambda^2;q^2)d_q y. \tag{3.26}$$

Proof. The first result follows immediately by proposition 3.6.
proving the second relation, by theorem 3.5 we have

$$\begin{aligned}
\varphi(x,\lambda^2;q^2) \quad &= \quad q^{-1}\sin(q\alpha;q^2)\cos(\lambda x;q^2) - q^{-\frac{1}{2}}\frac{\cos(q\alpha;q^2)}{\lambda}\sin(q^{\frac{1}{2}}\lambda x;q^2) \\
&+ \quad \frac{1}{\lambda}\int_0^\infty G(x,y,\lambda^2;q^2)p(y)\varphi(y,\lambda^2;q^2)d_q y - \frac{1}{\lambda}\int_x^\infty G(x,y,\lambda^2;q^2)p(y)\varphi(y,\lambda^2;q^2)d_q y
\end{aligned}$$

taking account of the fact that p in $L_q^1(\mathbb{R}_{q,+})$ and by (2.26) , (2.27)

$$\begin{aligned}
\mid \int_x^\infty G(x,y,\lambda^2;q^2)p(y)\varphi(y,\lambda^2;q^2)d_qy \mid &\leq C_q \int_x^\infty \mid p(y)\varphi(y,\lambda^2;q^2) \mid d_qy \\
&\leq C_q \parallel \varphi \parallel_{q,\infty} \int_0^\infty \mid p(y) \mid d_qy \\
&< +\infty.
\end{aligned}$$

leads to the result. □

On the same way, if we note by $\theta(x,\lambda^2;q^2)$ the solution of the q-difference problem

$$(E_2) = \begin{cases} L_qU = -\lambda^2 U \\ U(0,\lambda^2;q^2) = q^{\frac{1}{2}}\cos(q^{\frac{1}{2}}\alpha;q^2) \\ D_qU(0,\lambda^2;q^2) = -\sin(q^{\frac{3}{2}}\alpha;q^2) \quad ,\alpha \in \mathbb{C} \end{cases} \tag{3.27}$$

We show that

$$\theta(x,\lambda^2;q^2) = \mu_1(\lambda^2;q^2)\cos(\lambda x;q^2) + \nu_1(\lambda^2;q^2)q^{-\frac{1}{2}}\sin(q^{\frac{1}{2}}\lambda x;q^2) + \mathcal{O}(1;q^2) \tag{3.28}$$

where

$$\mu_1(\lambda^2;q^2) = q^{\frac{1}{2}}\cos(q^{\frac{1}{2}}\alpha;q^2) - \frac{1}{\lambda}\int_0^\infty sin(q^{\frac{3}{2}}\lambda y;q^2)p(y)\varphi(y,\lambda^2;q^2)d_qy \tag{3.29}$$

and

$$\nu_1(\lambda^2;q^2) = -q^{\frac{1}{2}}\frac{\cos(q\alpha;q^2)}{\lambda} - q^{\frac{1}{2}}\frac{\sin(q^{\frac{3}{2}}\alpha;q^2)}{\lambda}\int_0^\infty \cos(q\lambda y;q^2)p(y)\varphi(y,\lambda^2;q^2)d_qy. \tag{3.30}$$

Theorem 3.8. *For λ in $\mathbb{R}_{q,+}$, we have:*

$$\mu(\lambda^2;q^2)\nu_1(\lambda^2;q^2) - \nu(\lambda^2;q^2)\mu_1(\lambda^2;q^2) = \frac{1}{q^{\frac{1}{2}}\lambda}. \tag{3.31}$$

Proof. Using theorem 3.7, we obtain that

$$\varphi(x,\lambda^2;q^2) = \mu(\lambda^2;q^2)\cos(\lambda x;q^2) + \nu(\lambda^2;q^2)q^{-\frac{1}{2}}\sin(q^{\frac{1}{2}}\lambda x;q^2) + \mathcal{O}(1;q^2)$$

and

$$\theta(x,\lambda^2;q^2) = \mu_1(\lambda^2;q^2)\cos(\lambda x;q^2) + \nu_1(\lambda^2;q^2)q^{-\frac{1}{2}}\sin(q^{\frac{1}{2}}\lambda x;q^2) + \mathcal{O}(1;q^2).$$

We can deduce that

$$D_{q,x}\varphi(x,\lambda^2;q^2) = -q^{-1}\lambda\mu(\lambda^2;q^2)\sin(q\lambda x;q^2) + \lambda\nu(\lambda^2;q^2)\cos(q^{\frac{1}{2}}\lambda x;q^2) + \mathcal{O}(1;q^2)$$

and

$$D_{q,x}\theta(x,\lambda^2;q^2) = -\lambda q^{-1}\mu_1(\lambda^2;q^2)\sin(q\lambda x;q^2) + \lambda\nu_1(\lambda^2;q^2)\cos(q^{\frac{1}{2}}\lambda x;q^2) + \mathcal{O}(1;q^2),$$

then, the use of definition 3.3 leads to

$$\begin{aligned}
[\varphi,\theta]_q(x) &= \varphi(qx;q^2)D_{q,x}\theta(x;q^2) - \theta(qx;q^2)D_{q,x}\varphi(x;q^2) \\
&= \lambda\big[\nu(\lambda^2;q^2)\mu_1(\lambda^2;q^2) - \mu(\lambda^2;q^2)\nu_1(\lambda^2;q^2)\big]\big[\cos(q\lambda x;q^2)\cos(q^{\frac{1}{2}}\lambda x;q^2) \\
&+ q^{-\frac{3}{2}}\sin(q^{\frac{3}{2}}\lambda x;q^2)\sin(q\lambda x;q^2)\big] + \mathcal{O}(1;q^2).
\end{aligned}$$

In the other side, the fact that φ and θ are solutions of (E) and by proposition 3.4 we can deduce that

$$D_{q,x}\left[\varphi,\theta\right]_q(x)=0.$$

then from proposition 2.1

$$\begin{aligned}\left[\varphi,\theta\right]_q(x) &= \left[\varphi,\theta\right]_q(0)=q^{\frac{1}{2}}\cos(q^{\frac{1}{2}}\alpha;q^2)\cos(q\alpha;q^2)+q^{-1}\sin(q\alpha;q^2)\sin(q^{\frac{3}{2}}\alpha;q^2)\\ &= q^{\frac{1}{2}}\end{aligned}$$

the result follows. □

4. Asymptotic behaviour of $j_\alpha(\lambda x;q^2)$:

In this section, our objective is to establish; using the method of variation of constant given in the last section; the asymptotic expansion of $j_\alpha(\lambda x;q^2)$ when $\lambda\longrightarrow+\infty$.

we recall some properties given in [5]: For $\alpha>-\frac{1}{2}$, the q-Bessel function is defined by:

$$j_\alpha(x;q^2)=\Gamma_{q^2}(\alpha+1)\frac{q^\alpha(1+q)^\alpha}{x^\alpha}J_\alpha((1-q)x;q^2) \tag{4.1}$$

where $J_\alpha(x;q^2)$ is the q-Bessel Han Exton [15], defined by

$$J_\alpha(z;q)=\left(\frac{z}{1-q}\right)^\alpha\sum_{k=0}^{\infty}\frac{(-1)^kq^{\frac{k(k-1)}{2}}q^k}{\Gamma_q(k+1)\Gamma_q(\alpha+k+1)}\left(\frac{z}{1-q}\right)^{2k}. \tag{4.2}$$

This function satisfies the following relations

$$j_{-\frac{1}{2}}(x;q^2) = \cos(x;q^2) \tag{4.3}$$

$$j_{\frac{1}{2}}(x;q^2) = \frac{1}{x}\sin(x;q^2). \tag{4.4}$$

Proposition 4.1. *The function $j_\alpha(\lambda x;q^2)$; λ being complex; is the solution of the following q-difference problem:*

$$(E)=\begin{cases}\Delta_{q,\alpha}y(x)+\lambda^2y(x)=0\\ y(0)=1\\ D_qy(0)=0\end{cases} \tag{4.5}$$

where $\Delta_{q,\alpha}$ is the q-Bessel operator, defined by

$$\Delta_{q,\alpha}f(x) = \frac{1}{x^{2\alpha+1}}D_{q,x}\left[x^{2\alpha+1}D_{q,x}f\right](q^{-1}x), \tag{4.6}$$

$$= q^{2\alpha+1}\Delta_{q,x}f(x)+\frac{1-q^{2\alpha+1}}{(1-q)q^{-1}x}D_{q,x}f(q^{-1}x) \tag{4.7}$$

and

$$\Delta_{q,x}f(x)=(D^2_{q,x}f)(q^{-1}x) \tag{4.8}$$

Corollary 4.2. *For $x\in\mathbb{R}_q$ and $\frac{\log(1-q)}{\log q}\in\mathbb{Z}$, we have the following estimation*

$$|j_\alpha(x;q^2)| \leq \frac{1}{(q;q^2)^2_\infty} \tag{4.9}$$

Now, we consider the following q-Bessel equation given by:

$$(E)\qquad D^2_{q,x}y(x,\lambda^2;q^2)+\frac{\lambda^2}{q^{2\alpha+1}}y(qx,\lambda^2;q^2)=-\frac{1-q^{2\alpha+1}}{q^{2\alpha+1}(1-q)x}D_{q,x}y(x,\lambda^2;q^2).$$

Let $y_h(x,\lambda^2;q^2)=\sum_{n\geq 0}a_n(\alpha,\lambda^2;q^2)x^n$ the homogeneous solution of (E_h) given by

$$(E_h)\quad D^2_{q,x}y(x,\lambda^2;q^2)+\frac{\lambda^2}{q^{2\alpha+1}}y(qx,\lambda^2;q^2)=0.$$

Then, we have

$$D^2_{q,x}y(x,\lambda^2;q^2)=\sum_{n\geq 2}\frac{(1-q^n)(1-q^{n-1})}{(1-q)^2}a_nx^{n-2}=\sum_{n\geq 0}\frac{(1-q^{n+2})(1-q^{n+1})}{(1-q)^2}a_{n+2}x^n.$$

By identification, we obtain the following recurrence relation:

$$\frac{(1-q^{n+2})(1-q^{n+1})}{(1-q)^2}a_{n+2}=-\frac{\lambda^2q^n}{q^{2\alpha+1}}a_n. \tag{4.10}$$

If we proceed in a similar way of proposition 2.3, we can deduce easily that the homogeneous solution $y_h(x,\lambda^2;q^2)$ is given by

$$y_h(x,\lambda^2;q^2)=a_0\cos(q^{-\alpha-\frac{1}{2}}\lambda x;q^2)+a_1\frac{q^{-\alpha-1}}{\lambda}\sin(q^{-\alpha}\lambda x;q^2) \tag{4.11}$$

where a_0,a_1 are constants in $\mathbb{R}$.
Now, we give a particular solution $y_p(x,\lambda^2;q^2)$ of (E) in the form

$$y_p(x,\lambda^2;q^2)=a_\alpha(x,\lambda^2;q^2)\cos(q^{-\alpha-\frac{1}{2}}\lambda x;q^2)+b_\alpha(x,\lambda^2;q^2)\frac{q^{-\alpha-1}}{\lambda}\sin(q^{-\alpha}\lambda x;q^2). \tag{4.12}$$

The q-wronskian is given by

$$\begin{aligned}W(x,\lambda^2;q^2)=&\ D_{q,x}\left[\cos(q^{-\alpha-\frac{1}{2}}\lambda x;q^2)\right]\frac{q^{-\alpha-1}}{\lambda}\sin(q^{-\alpha}\lambda x;q^2)\\&-\cos(q^{-\alpha-\frac{1}{2}}\lambda x;q^2)D_{q,x}\left[\frac{q^{-\alpha-1}}{\lambda}\sin(q^{-\alpha}\lambda x;q^2)\right],\end{aligned}$$

thus, using corollary 2.4 we can show that

$$D_{q,x}W(x,\lambda^2;q^2)=0,$$

therefore

$$W(x,\lambda^2;q^2)=-q^{-2\alpha-1}\lambda. \tag{4.13}$$

Using the q-method of variation of constant (given in theorem 3.5) and proposition 2.1 we deduce that

$$a_\alpha(x,\lambda^2;q^2)\ =\ \frac{q^{-\alpha-1}}{\lambda}\int_x^{+\infty}\frac{1-q^{2\alpha+1}}{1-q}\frac{1}{t}D_{q,t}y(t,\lambda^2;q^2)\sin(q^{-\alpha+1}\lambda t;q^2)d_qt \tag{4.14}$$

$$b_\alpha(x,\lambda^2;q^2)\ =\ -\frac{1}{\lambda}\int_x^{+\infty}\frac{1-q^{2\alpha+1}}{1-q}\frac{1}{t}D_{q,t}y(t,\lambda^2;q^2)\cos(q^{-\alpha+\frac{1}{2}}\lambda t;q^2)d_qt \tag{4.15}$$

this leads to the following result

Proposition 4.3. *For $x, \lambda \in \mathbb{R}_{q,+}$, the solution $y(x, \lambda^2; q^2)$ of (E) is given by:*

$$\begin{aligned} y(x,\lambda^2;q^2) &= a\cos(q^{-\alpha-\frac{1}{2}}\lambda x;q^2) + \frac{b}{\lambda}q^{-\alpha-1}\sin(q^{-\alpha}\lambda x;q^2) \\ &+ \frac{q^{-\alpha-1}}{\lambda}\int_x^{+\infty} G_\alpha(t,x,\lambda;q^2)\frac{1-q^{2\alpha+1}}{1-q}\frac{1}{t}D_{q,t}y(t,\lambda^2;q^2)d_qt \end{aligned}$$

where

(4.16)

$$G_\alpha(t,x,\lambda;q^2) = \cos(q^{-\alpha-\frac{1}{2}}\lambda x;q^2)\sin(q^{-\alpha+1}\lambda t) - \cos(q^{-\alpha+\frac{1}{2}}\lambda t;q^2)\sin(q^{-\alpha}\lambda x;q^2)$$

On the other hand $j_\alpha(\lambda x; q^2)$ is the unique solution of (E) with initial conditions

$$\begin{aligned} j_\alpha(0;q^2) &= 1 \\ D_q j_\alpha(0;q^2) &= 0. \end{aligned}$$

Therefore, we can write $j_\alpha(\lambda x; q^2)$ as the following form

Theorem 4.4. *For $\lambda \in \mathbb{R}_{q,+}$ and $\alpha > -\frac{1}{2}$, we have*

(1)

$$j_\alpha(\lambda x;q^2) = \cos(q^{-\alpha-\frac{1}{2}}\lambda x;q^2) + R_{q,\alpha}(x,\lambda^2), \tag{4.17}$$

where

$$R_{q,\alpha}(x,\lambda^2) = -\frac{1}{\lambda}\frac{1-q^{2\alpha+1}}{1-q}\int_x^{+\infty} G_\alpha(t,x,\lambda;q^2)\frac{1}{t}D_{q,t}j_\alpha(\lambda t;q^2)d_qt$$

(2) *Additionally, $R_{q,\alpha}(x, \lambda^2)$ tend to 0 when λ tend to $+\infty$.*

Proof. To prove 2. it suffices to use (2.26), (2.27) and corollary4.2, then

$$\begin{aligned} |R_{q,\alpha}(x,\lambda^2)| &\le \frac{1}{\lambda}\frac{1-q^{2\alpha+1}}{1-q}\frac{2}{(q;q^2)^2_\infty}\int_x^{+\infty} \mid \frac{D_{q,t}j_\alpha(\lambda t;q^2)}{t} \mid d_qt \\ &\le \frac{1-q^{2\alpha+1}}{1-q}\frac{2}{(q;q^2)^2_\infty}\int_{\lambda x}^{+\infty} \mid \frac{D_{q,t}j_\alpha(t;q^2)}{t} \mid d_qt \\ &\le \frac{1-q^{2\alpha+1}}{1-q}\frac{2}{(q;q^2)^2_\infty}\int_{\lambda x}^{+\infty} \frac{|j_\alpha(t;q^2)-j_\alpha(qt;q^2)|}{(1-q)t^2}d_qt \\ &\le \frac{1-q^{2\alpha+1}}{(1-q)^2}\Big[\frac{2}{(q;q^2)^2_\infty}\Big]^2\int_{\lambda x}^{+\infty}\frac{d_qt}{t^2} \\ &\le \frac{C_q}{\lambda x} \longrightarrow 0 \quad ,\lambda \longrightarrow \infty \end{aligned}$$

where

$$C_q = \frac{1-q^{2\alpha+1}}{q}\Big[\frac{2}{(1-q)(q;q^2)^2_\infty}\Big] \tag{4.18}$$

and the result follows immediately. □

4.1. **Application.** We recall the q-equality of Weber integral study in [5]. For a, λ in $\mathbb{R}_{q,+}$ and $\alpha > -1$ we have

$$\frac{1}{A_\alpha}\int_0^\infty e(-a^2x^2;q^2)j_\alpha(\lambda x;q^2)x^{2\alpha+1}d_qx = \frac{1}{a^{2\alpha+2}}e(-\frac{\lambda^2}{a^2(1+q)^2};q^2) \tag{4.19}$$

where

$$A_\alpha = \int_0^\infty \frac{x^{2\alpha+1}}{(-(1-q^2)x^2;q^2)} d_q x \tag{4.20}$$

and if we take $a = \sqrt{t}$ we obtain

$$E_\alpha(t,\lambda^2;q^2) = \int_0^\infty e(-q^{-1}tx^2;q^2) j_\alpha(\lambda x;q^2) x^{2\alpha+1} d_q x \tag{4.21}$$

where

$$E_\alpha(t,\lambda^2;q^2) = \frac{A_\alpha}{t^{\alpha+1}} e(-\frac{\lambda^2}{(1+q)^2 t};q^2) \tag{4.22}$$

Proposition 4.5. *For $\alpha > -1$ and $\lambda, t \in \mathbb{R}_{q,+}$, the heat kernel $E_\alpha(t,\lambda;q^2)$ has the following behaviour:*

$$E_\alpha(t,\lambda^2;q^2) = \int_0^\infty e(-q^{-1}tx^2;q^2)\cos(q^{-\alpha-\frac{1}{2}}\lambda x;q^2) x^{2\alpha+1} d_q x + \Theta_\alpha(\lambda,t;q^2) \tag{4.23}$$

where

$$\Theta_\alpha(\lambda,t;q^2) \longrightarrow 0 \quad ,\lambda \to \infty \tag{4.24}$$

Proof. the result follows by proposition 4.4 and the fact that

$$\begin{aligned} | \int_0^\infty e(-q^{-1}tx^2;q^2) R_q(\lambda^2,x) x^{2\alpha+1} d_q x \, | &\leq \frac{C_q}{\lambda^2} \int_0^\infty e(-q^{-1}tx^2;q^2) x^{2\alpha} d_q x \\ &= \frac{C_q}{\lambda^2}(1-q) \sum_{-\infty}^{+\infty} \frac{q^{(2\alpha+1)k}}{(-q^{-1}(1-q^2)tq^{2k};q^2)_\infty} \\ &= \frac{C_q}{\lambda^2}(1-q) \sum_{-\infty}^{+\infty} \frac{q^{2\beta k}}{(aq^{2k};q^2)_\infty} \end{aligned}$$

where $\beta = \alpha + \frac{1}{2}$ and $a = -q^{-1}(1-q^2)t$. The use of Ramanujan's sum (see [7]) leads to

$$| \int_0^\infty e(-q^{-1}tx^2;q^2) R_q(\lambda^2,x) x^{2\alpha+1} d_q x \, | \leq \frac{C_q}{\lambda^2} B_\alpha(t;q^2) \longrightarrow 0 \quad ,\lambda \to \infty$$

where

$$B_\alpha(t;q^2) = \frac{(-q^{2\alpha}t(1-q^2), q^{2-\alpha}t^{-1}(1-q^2)^{-1}, q^2;q^2)_\infty}{(q^{2\alpha+1}, -q^{-1}t(1-q^2), -q^3t^{-1}(1-q^2)^{-1};q^2)_\infty}. \tag{4.25}$$

□

References

[1] **R.Askey, N.Atakisheiev and S.Suslov**.*An analog of the Fourier transformation for the q-harmonic oscillator*, Symmetries in Science VI(1993)57-63(B.Gruber, ed)Plenum Press, NY.

[2] **Agranovich and Marchenko**, *the inverse problem of scatering theory* , Gordan and Breach , New York 1963.

[3] **PH.Feinsilver**,*Elements of q-Harmonic Analysis*, J.Math.Anal.141, 1989, 509-526.

[4] **A.Fitouhi, and F.Bouzeffour**, *q-Cosine Fourier Transform and q-Heat Equation* , to appear in Ramanujan Journal.

[5] **A.Fitouhi, M.Hamza and F.Bouzeffour**, *The q-j_α Bessel function*, journal of approximation Theory 115, 114-116(2002).

[6] **G.Gasper and M.Rahmen**, *Basic Hypergeometric series* , Encyclopedia of mathematics and its applications 35 , Cambridge university press, 1990.

[7] **M.E.H.Ismail**, *A simple proof of Ramanujan's sum* , proc.Amer.Math.Soc.63 .1977, 185-186.

[8] **F.H.Jackson** , *On q-Functions and a certain Difference Operator* , Transactions Of the Royal Society of London 46, 1908, 253-281.

[9] **F.H.Jackson**, *On a q-Definite Integrals,* Quarterly Journal of Pure and Application Mathemathics 41, 1910, 193-203.

[10] **T.H.Koornwinder**, *q-Special Function*, a Tutorial, Mathematecal Preprint Series, report 94-08, Univer.Amesterdam.Th Netherlands.

[11] **T.H.Koornwinder and R.F.Swarttouw :***Special Function,and q-commuting variable, in Special Function,q-Series and Related Topics,* M.E.Ismail,D.R.Masson and Rahman(eds), Fields Institue Communication 14, American Mathematical Society,1997,pp.131-166.

[12] **T.H.Koornwinder and R.F.Swartouw**, *On q-Analogues of the Hankel and Fourier Transform,* Tran.AM.S.1992, 333, 445-461.

[13] **A.Kempf and S.Majid**, *Algebraic q-Integration and Fourier Theory on Quantum and Braided Spaces,* J.Maths.Phys.35 (1994) 6802-6837.

[14] **V.A.Marchenko**, *Some questions in the theory of one-dimential linear differential operators of the second order,* Amer.Math.Soc. Transl. vol 101, 1973.

[15] **R.F.Swarttouw**, *The Han-Exton q-Bessel Function,* Thesis.

[16] **Ian N.Sneddon**, *The Use of Integral Transform,* McGraw-Hill Book Company.

[17] **G.N.Watson**, *A Treatise on the Theory of the Bessel Functions,* Lnd ed.Combridge Uni.Press.London and New-York.

Département de Mathematiques, Faculté des Sciences de Tunis, 1060 Tunis, Tunisia.
E-mail address: `Ahmed.Fitouhi@fst.rnu.tn`

Département de Mathematiques, Faculté des Sciences de Tunis, 1060 Tunis, Tunisia
E-mail address: `Akram.Nemri@fst.rnu.tn`

Faculté des Sciences de Tunis, 1060 Tunis, Tunisia.
E-mail address: `Meniar.Haddad@fst.rnu.tn`